
AGRICULTURE IN VANUATU PLANT BIOLOGY




NOTES FOR TEACHERS



Agriculture in Vanuatu

PLANT BIOLOGY

(31)



Ministry of Education Port Vila Republic of Vanuatu 1997



Revised Edition 1997

This book was written by S. J. Goodyer and first published in 1987.

This revised edition has been prepared by the Ministry of Education under the auspices of the Primary and Secondary Education Project with support from the World Bank (Credit 1964-VAN).

© Ministry of Education

All rights reserved. No part of this publication may be reproduced, stored in a retrieval system or transmitted, in any form or by any means, without prior permission from the publisher.


Contents

1.	Parts of a plant	5
2.	Leaves	7
3.	Transpiration	9
4.	Photosynthesis	12
5.	Respiration	15
6.	Stems	17
7.	Roots	21
8.	Flowers	24
9.	Pollination	26
10.	Fertilisation	27
11.	Fruits	28
12.	Seeds	30
13.	Germination	32
	Glossary	34

1. Parts of a plant

A plant is made up of many parts or structures. In order to understand how a plant works we must learn about these different parts.

Figure 1 Parts of a plant



Shoot

the part of the plant growing above the soil. It consists of stems, plus buds and leaves.

Main shoot

the first shoot to grow, and usually the biggest.

Side (lateral) shoots

These grow later from the **axils** of the **main shoot**. They are usually smaller and sometimes called **branches**.

Axil

the gap between a leaf and the stem. Axillary

buds are found here.

Mode

the position where a leaf joins the stem. The

lowest node is the 1st node, next one up the

2nd node, and so on.

Internode

the gap between two nodes.

Buds

these produce new cells to make the plant grow. They can produce **shoots**, **flowers**, or **fruits**. They are sometimes called the **growing**

points.

Terminal bud

this is found at the top of the main shoot. It

makes the plant grow tall and straight.

Axillary buds

these occur in the axils. They sometimes make

the plant grow "bushy" by producing side

shoots.

Leaf scar

a mark on the stem showing from where a leaf

has fallen.

Practical work

After the introduction of this topic, students should make a collection of different plants from the garden or bush. They should try to identify the various structures present on these plants.

2. Leaves

Function

The main function of leaves is to catch **sunlight**, and make plant food by **photosynthesis**.

Types


There are two main types of leaf:

1. Narrow Leaves

These have straight parallel veins and a leaf sheath.

Plants with leaves like this are called monocots.


For example: corn and grasses



2. Broad Leaves

These have branching veins, sometimes called **net veins**, and they have a **petiole** (**leaf stalk**). Plants with leaves like this are called **dicots**.

For example: taro and pawpaw



Leaf veins carry water, nutrients and plant food through

the leaf. The main vein in the centre of the leaf is

called the midrib.

Leaf blade is the main part of the leaf structure. It is flat

and thin to allow it to catch plenty of sunlight.

The top of the leaf blade is **dark green**, this is because it contains a substance called **chlorophyll**. Chlorophyll helps the leaf absorb sunlight energy.

The **bottom** of the leaf blade is full of small holes called **stomata**. These are too small for us to see, but very important, because they allow **water** and **gases** to move in and out of the leaf.

Practical work

Students should make a collection of different leaves from the garden or bush. They should try to classify them as Broad or Narrow leaves and identify the various parts.

3. Transpiration


This is the loss of water from plants. It occurs mainly in the leaves.

The transpiration process

- (i) Sunlight falls onto plant leaves.
- (ii) The heat of the sun turns the water inside the leaf cells into water vapour. The water vapour moves from the cells into the air spaces.
- (iii) The water vapour moves out of the leaves through the **stomata** and is lost to the air by **evaporation**.

Figure 4

Transpiration process in a leaf



A small amount of water is lost from plant stems through tiny holes called **lenticels**.

Transpiration helps to keep leaves cool (like sweating in animals).

Conditions which speed up transpiration

The following weather conditions accelerate the loss of water from leaves:

1. Sunlight

because sunlight heats up the water inside leaves.

2. Wind

because wind increases the rate of evaporation from stomata.

3. Dry Atmosphere

because it is easier for water to evaporate into dry air than humid air. This is because humid air already contains a lot of moisture.

In these weather conditions it is therefore very important to keep crop plants supplied with plenty of water. If plants become short of water, they wilt and die.

Practical work


The following is a very simple demonstration of transpiration.

Procedure

- (i) Take two, clear, dry plastic bags.
- (ii) Place one of these over a small branch of a plant growing in the sun and tie the mouth of the bag tightly around the stem (see Figure 5).
- (iii) Place the second bag over a similar branch which has had all its leaves removed (see Figure 5).
- (iv) After 2 hours return to the plant and examine the bags carefully.
- (v) Observe the presence of water on the inside of the plastic bag covering the leafy branch, whilst the bag covering the leafless branch is dry.

Explain to students that this demonstrates that water is lost from the leaves.

Transpiration Demonstration



4. Photosynthesis

This is the name given to the process by which plant leaves use sunlight energy to make simple foods.

These simple foods are called **carbohydrates**, and they are rich in **energy**.

The carbohydrates we find in plants are sugars and starch.

The photosynthesis process


- (i) Sunlight energy is absorbed by chlorophyll at the top of the leaf.
- (ii) Carbon dioxide gas enters the leaf through the stomata at the bottom.
- (iii) Water enters the leaf through the veins. (It enters the plant through the roots.)

Then sunlight, carbon dioxide and water react together to make carbohydrate (sugar) and oxygen.

The photosynthesis equation

Figure 6

Photosynthesis in the leaf



Oxygen gas produced by photosynthesis moves out into the air. through the Stomata.


Sugars produced by photosynthesis can be used in two ways:

- To give plants energy to grow (see notes on respiration).
- They can be moved to another part of the plant (usually the root or stem) and changed into starch. This happens in many of our food crops.

e.g. yam, taro, manioc and kumala starch acts as a store of energy.

Figure 7

Starch storage in taro



Practical work


...nstrate that Students can carry out the following experiment to demonstrate that light is needed for photosynthesis.

Procedure

- (i) Cover the leaf of a tree that is growing in sunlight with a strip of silver paper (see Diagram 8).
- (ii) After 3 days remove the leaf from the tree, take off the silver paper and kill the leaf by boiling it in water.
- (iii) Remove the chlorophyll from the leaf by washing it with methylated spirits.
- (iv) Place the leaf in iodine solution and observe how the part of the leaf not covered by silver paper turns a blue-black colour.

Explain to the students that iodine only turns blue-black in the presence of starch. The reason why the part of the leaf under the silver paper does not turn blue-black is because there was no sunlight for it to make starch by photosynthesis.

Figure 8 To show that light is required for photosynthesis



5. Respiration

This is the process which all plants (and animals) use to obtain energy from simple food (carbohydrate).

Energy is required by plants for growth, movement and other processes which take place inside them.

The respiration process

Respiration takes place inside the living plant cells. For respiration to occur, the cells require:


- (i) Carbohydrate: this is made by plant leaves during photosynthesis.
- (ii) Oxygen: this is taken in from the air through the stomata.

During the respiration process the carbohydrate is broken down to release the energy stored inside it.

The respiration equation:

The carbon dioxide gas is released back into the air through the stomata.

- N.B. Photosynthesis and respiration are "reverse processes":
 - during **photosynthesis**, energy from the sun is **taken in** by plants and **stored** in simple foods.
 - during **respiration** that energy which was taken in is **released** from the simple foods, and used by the plants for growth, movement etc.



Practical work


It is not easy to demonstrate respiration in plants. After explaining that plants breathe out the same gases as humans, the following limewater test for carbon dioxide can be carried out:

Procedure

- (i) Heat Calcium Carbonate (coral) strongly to change it to quicklime (Calcium Oxide).
- (ii) Add the quicklime to water, shake and stand for a few days in a covered container until the top of the liquid is clear. If possible, filter it.
- (iii) Add some of the clear liquid to a test tube or small bottle and breathe into it using a straw or thin tube (see Figure 9).
- (iv) Observe the clear liquid turn a milky-white colour.

Explain to the students that this change occurs only in the presence of carbon dioxide.

Figure 9 Limewater test for carbon dioxide



6. Stems

Functions

Stems have three main functions:

1. Support


They hold up other parts of the plant that are above the soil. eg. leaves, flowers and fruits.

2. Transport

- (i) Water and nutrients move upwards through the centre of the stem.
- (ii) Sugars move downwards, from the leaves, through the outside of the stem.

Figure 10

Transportation in a stem



Key

+++-->--- sugars from the leaves

----- water and nutrients from the roots

3. Food Storage


Most juicy stems store some food. Some have special forms for this purpose. The food is usually stored as **starch**. Starch is a **carbohydrate** made during photosynthesis and is rich in **energy**. Here are some examples:

(i) Stem Tubers

These are swollen underground stems. eg. Yam

Figure 11

Stem tuber




(ii) Rhizomes

These are swollen underground stems which are horizontal. e.g. Ginger

Figure 12

Rhizome




(iii) Corms

This is a swollen stem base. eg. Taro

Figure 13






Stolons

These are horizontal stems which grow just above the soil. Usually they do not store food, but help the plant to spread out and cover the round, eg. buffalo grass, watermelon.

Figure 14

Stolon



Practical work

- 1. Students should make a collection of the different kinds of stems outlined in the notes, and draw labelled diagrams.
- 2. The following simple experiment can be used to demonstrate the pathway of water through the stem:

Procedure

- (i) Pick several small leafy plants and cut across the bottom end of their stems.
- (ii) Place the cut ends of the stems in a coloured solution such as neutral red or methylene blue.
- (iii) After five minutes remove one of the plants and slice across the bottom of the stem. Observe the water vessels which will have been stained by the coloured solution.
- (iv) Cut sections higher and higher up the stem until the colour can no longer be seen. This will show the height to which the solution has risen in that time.
- (v) Remove another plant after ten minutes, then another after fifteen minutes and so on repeating the procedure (iv). This will show the rate at which the solution is rising up the stem.

7. Roots

Functions

Roots have 2 main functions:


1. Anchorage:

They hold plants firmly in the soil.


2. Absorption:

They absorb water and nutrients from the soil. This is done by the root hairs. These are found at the end of the main roots. These are small and easily damaged.

Figure 15 Position of root hairs



The root hairs absorb water and nutrients from the layer of water which surrounds the soil particles.



Root systems

There are two main types of root systems:-

(i) Tap Root System


This system has one main root which can grow quite deep, and is often thick. Other smaller roots called **lateral roots** branch from it (see Figure 17).

(ii) Fibrous Root System

This system has roots which all grow to about the same size and length. This system is usually not as deep as a tap root system (see Figure 17).

Figure 17

Types of root system



Food storage in roots

Some plants use their roots to store food. The food is usually stored as starch. Starch is a carbohydrate, rich in energy.


The two main types of root structures which store food are:

(i) Swollen Tap Roots

The tap root of the plant becomes swollen with starch. eg. carrot, beetroot, radish.

Figure 18

Tap root of a radish plant




(ii) Root tubers

Parts of the root system become swollen with starch, eg. kumala, manioc.



Tuber roots of kumala



Practical work

Students should make a collection of different plants to discover if they have Tap Roots or Fibrous Roots, and observe the Root Hairs. They should also observe as many as possible of the other root features outlined in the notes.


8. Flowers

Function

The function of flowers is to help the plant to reproduce by making seeds. These are usually contained in a **fruit**.

Flowers can be many different colours, shapes and sizes. Most contain male and female parts but some, for example pawpaw and corn, have separate male and female flowers.

Figure 20 Parts of an insect pollinated flower



Parts of the flower

(see Figure 20)

Flower part		function	
Γ	sepals	protect the flower bud.	
	stalk	supports the flower, and transports water and nutrients to it.	
	petals	often brightly coloured to attract insects.	
Female parts (Capel)	stigma	collects pollen from insects or the air.	
	stylę	supports the stigma, and pollen grows down it.	
	ovary	contains ovules, and becomes the fruit.	
\ L	ovules	become the seeds.	
Male parts	— anther	produces pollen.	
(Stamen)	men) filament	supports the anther.	
	nectary insects (not p	sweet sugary substance. It attracts present in all flowers).	

Practical work

Students should make a collection of flowers from the garden or bush, and attempt to identify the various structures in tomato, pigeon pea, Christmas tree, funeral lily. Using a razor blade and magnifying glass, the flowers can be cut in half to observe the ovules.

Many flowers have difficult structures. It is probably best to avoid bougainvillea, hibiscus, frangipani, pawpaw and coconut.


9. Pollination

This is when pollen moves from an anther to a stigma.


Self pollination is when pollen moves onto the stigma of the **same** flower.

Cross pollination is when pollen moves onto the stigma of a different flower.

Figure 21



Pollination



Self pollination

Cross pollination

How pollen moves

Pollen is moved from an anther to a stigma by insects or wind:

1. Insect Pollination

Insects are attracted to flowers with bright coloured petals and sweet smelling scents. The insects pick up pollen on their bodies as they search for nectar. This pollen can then later be brushed onto a stigma, causing pollination.

2. Wind Pollination

This occurs in flowers which are usually small and have no bright coloured petals or scent. Such flowers, for example grasses, produce a lot of light pollen which can be blown from an anther to a stigma by the wind.

Practical work

Students should be asked to find 3 examples of insect-pollinated flowers, and 3 wind-pollinated flowers.

10. Fertilisation

After pollination the pollen grains on the stigma produce a pollen tube. Each pollen tube then grows down the style, until it reaches the ovary. Once inside the ovary, the pollen tube will join with an ovule, this is called fertilisation.

After fertilisation the following changes take place:

Sepals usually die and drop off.

Petals usually die and drop off.

Stigma and style usually die and drop off


Anther and filament usually die and drop off.

Ovules become the seeds

Ovary becomes the fruit

Figure 22

Fertilisation of the flower



Practical work flower

stalk Collect some tomato flowers and green tomato fruits. Using razor fruit, stal blades, students can dissect the flowers and fruits to observe the stages of fruit formation.

11. Fruits

A fruit is a swollen ovary containing one or more seeds.

16 per 25 cores sueurs

Functions

The two main functions of fruits are:

- 1. Protection of the seeds
- 2. Dispersal of (scatter) the seeds.

Seed disposal

There are several methods used by different fruits to disperse seed:

(i) Wind Dispersal

The fruits are small and light so that they can be carried by the wind. Many weeds are dispersed this way.

(ii) Animal Dispersal

Succulent fruits are often eaten by birds or mammals. The seeds of these are either thrown away or they pass through the digestive system to be dispersed in the faeces, eg. tomato.

(iii) Water Dispersal


Seeds which float on water can be dispersed by river or sea, eg. coconuts.

(iv) Self Dispersal

The fruit wall dries out creating a tension. It then bursts open suddenly throwing the seeds out.


Figure 23

Kinds of fruit



Practical work

Students should make a collection of as many different kinds of fruits as possible. They should dissect them to observe the various features outlined in the notes. For each fruit they should attempt to determine the method of dispersal used.



12. Seeds

A seed develops from a fertilised ovule.

Function

The function of all seeds is to find new ground and grow into plants.

Types

Seeds are grouped into two types:-

1. Monocots

These have 1 **cotyledon**, for example: most narrowleaved plants, such as corn, grass.

2. Dicots

These have 2 cotyledons, for example: most broadleaved plants, such as taro and pawpaw.

Parts^{*}

A seed has three main parts:

1. Testa

This is the seed coat. It protects the soft parts inside.

2. Embryo

This will grow into the new plant after the seed germinates. It consists of two parts:

(i) radicle: the young root

(ii) plumule: the young shoot


3. Food Store


This supplies energy to the embryo when it begins to grow. **Dicots** store food in the **cotyledons**.

Monocots store food in the endosperm.

Figure 24

Parts of seeds





Scar:

place where seed was joined to fruit.

Micropyle:

hole in testa through which water and air are

taken in during germination.

Practical work

Take a sample of corn and bean seeds. Get the students to soak these in water for four or five days. After this time they will be able to dissect the seed to observe the various features inside.

13. Germination

This is when the seed begins to grow into a plant.

Conditions for germination

Some seeds can remain alive in the soil for many years. For seeds to germinate they need to have enough:

- water
- air
- warmth

Note: Seeds do not require light to germinate and most seeds will germinate in light or dark conditions.

Stages of germination

- 1. The seed takes in air and water which causes it to swell up.
- 2. The testa splits open and the **radicle** emerges. The radicle grows down into the soil.
- 3. The plumule emerges and grows upwards towards the light.


The energy for germination is supplied by the food store inside the seed (cotyledons or endosperm).

As soon as the young plant has developed leaves it can make its own food by **photosynthesis**.

The young plant is called a seedling (see Figure 25).

Practical work

Prior to teaching this topic plant some bean seeds out so that you have seeds at various stages of germination available for observation by the students. Start with the earliest planting at about two weeks before, and the last at about 4 days before teaching.



Glossary

carbohydrate simple food, made by plants during

photosynthesis, which are rich in energy

(eg. sugars and starch).

carbon dioxide a gas, present in the air, taken in by plants

during photosynthesis, and given out

during respiration.

chlorophyll a green chemical which absorbs sunlight.

corm swollen stem base which stores food.

cotyledon a seed leaf.

cross pollination pollination occuring between different

flowers.

dicot plant which has two cotyledons in its seed.

embryo the part of the seed which will grow into a

plant.

endosperm food store found in some seeds.

fertilisation when pollen joins with an ovule.

fibrous roots a root system where all the roots of a

plant grow to approximately the same

size.

germination when a seed begins to grow into a plant.

insect pollination pollination carried out by insects.

leaf blade the flat part of a leaf which catches

sunlight for photosynthesis.

leaf veins tubes running through the leaf which carry

water, nutrients, and plant food.

lenticels

small holes found on outside of plant

stems.

micropyle

small hole in the seed coat (testa) through

which water and air are taken in.

midrib

the main vein running through the middle

of a leaf.

monocot

plant which has one cotyledon in its seed.

net veins

a system of branching veins found in

broad leaves.

oxygen

gas, present in the air, taken in by plants

for respiration and given out after

photosynthesis.

petiole

leaf stalk found in broad leaved plants

which attaches leaf blade to stem.

photosynthesis

the production of food by green plants.

plumule

the shoot of a seed embryo.

pod

fruit belonging to a legume plant.

pollination

when pollen moves from an anther to a

stigma.

radicle

the root of a seed embryo.

respiration

process used by plants (and animals) to

obtain energy from simple foods.

rhizone

horizontal underground stem.

root hair

part of the root which absorbs water and

nutrients from the soil.

root tuber

swollen root containing stored food.

seedling

the young plant which emerges after

germination.

self pollination pollination occuring within a flower.

stem tuber underground stems, swollen with food.

stolon horizontal stem running overground.

stomata small holes in bottom of leafthrough which

water and gases can pass.

swollen tap root tap root which becomes swollen with

food.

tap root root, present in some plants, which can

grow deep and become quite thick. Other

smaller roots branch from it.

testa the seed coat or outside skin of the seed.

transpiration loss of water from plant shoots mainly

from evaporation from the leaves.

wind pollination pollination carried out by wind.

